MULTIMEDIA UNIVERSITY

STUDENT ID NO

MULTIMEDIA UNIVERSITY

FINAL EXAMINATION

TRIMESTER 2, 2015/2016

HEM 2019 PROBABILITY AND BIOSTATISTICS

(All sections / Groups)

11th MARCH 2016
9:00 – 11:00 AM
(2 Hours)

INSTRUCTIONS TO STUDENTS

1. This Question paper consists of 14 pages with 5 Questions only.
2. Attempt **FOUR** out of FIVE questions. All questions carry equal marks and the distribution of the marks for each question is given.
3. Please print all your answers in the Answer Booklet provided.
4. Please refer to the appendix for the formula and statistical tables.
5. You may use calculators in this examination.

QUESTION 1

a. Suppose that exposure to a particular chemical solvent commonly used in manufacturing increases the risk of bladder cancer; 13 of the workers who had bladder cancer were exposed to the solvent while 56 were not. Of the workers without bladder cancer, four had been exposed to the solvent while 65 had not.

		Exposed to solvent	
		Yes	No
Bladder Cancer	Yes	13	56
	No	4	65

Use this data to compute the sample odds ratio. Show all your works. [2 marks]

b. Suppose that the probability of developing lung cancer among smokers is 15% and among non-smokers is 2%, calculate the relative risk of developing lung cancer associated with smoking. Show all your works. [2 marks]

	Lung cancer	
	Yes	No
Smoker	15	85
Non-smoker	2	98

c. What are the two main difficulties in observational studies? [2 marks]

d. Why doesn't a statistically significant difference necessarily mean a biological significance? [2 marks]

e. Sampling and nonsampling errors may occur during data collection. Give two examples of nonsampling errors. [2 marks]

QUESTION 2

a. The probability that a person selected at random from a population will be tested positive for disease A is 0.1, and the probability that a person selected at random has the disease A is 0.13. The probability that a person who is tested positive also has the disease A is 0.08.

i. Compute a contingency table. [2 marks]

ii. What is the specificity of the test? [1 mark]

iii. A person selected at random from the population is tested negative. What is the probability that the person has the disease? [1 mark]
Show all your works.

Continued...

QUESTION 2 (Cont')

b. In a certain population of herring fish, the lengths of the individual fish follow a normal distribution. The mean length of the fish is 54.0 mm, and the standard deviation is 4.5 mm.

i. What percentage of the fish are less than 60 mm long? [2 marks]
ii. What percentage of the fish are more than 51 mm long? [2 marks]
iii. What is the maximum fish length for fishes in the bottom 20%? [2 marks]
Show all your works.

QUESTION 3

a. In Mendel's classical experiments with peas, he produced hybrids according to the ratio of 9: 3: 3: 1 for different phenotypes; round and yellow, round and green, angular and yellow, and angular and green respectively. In an experiment designed to test whether inheritance in peas follow the claimed ratio, the following results were recorded from 760 dihybrid crosses.

Type	Observed frequency
Round and yellow	378
Round and green	167
Angular and yellow	128
Angular and green	87

Do the observations show that peas are following the Mendel ratio? Test using the χ^2 goodness-of-fit test with $\alpha = 0.05$. Show all your works. [6 marks]

b. In the simplest form of Mendelian inheritance, a physical characteristic of a plant or an animal is determined by a single pair of genes. The color of seeds from sweet pea plants is an example. Let y and g represent yellow and green genes, seeds will be green if the plant has the color-gene pair (g, g); it will be yellow if the plant has color-gene pairs of (y, y) or (y, g). When two (y, g)-plants are crossed,

i. What are the possible genotypes and phenotypes that the cross can produce? [2 marks]
ii. What is the probability that a yellow seed will be the result? [1 mark]
iii. What is the probability that a yellow seed will be (yy)? [1 mark]
Show all your works.

Continued...

QUESTION 4

Measurement of serum cholesterol and arterial calcium deposition was made on 6 patients. The data are shown as below.

Calcium, mg/dl (X)	Cholesterol, mg/dl (Y)
59	298
52	303
43	232
59	288
24	235
37	267

i. Calculate the linear regression of Y on X. [4 marks]
ii. Plot the data and draw the regression line on your graph. [2 marks]
iii. Calculate the sample coefficient of determinations, r^2. [2 marks]
iv. Find the predicted value of cholesterol if calcium is 45 mg/dl. [2 marks]
Show all your works.

QUESTION 5

a. The diameter of the stem of a plant is an important trait because of its relationship to breakage of the stems, which interferes with harvesting the crop. The stem diameters were measured after flowering of the plant. The mean and standard deviation of stem diameters (mm) for 8 samples were 2.275 and 0.2375, respectively.
i. Calculate the standard error of the mean. [2 marks]
ii. Construct a 95% confidence interval for the population mean. [2 marks]
Show all your works.

b. A medical products firm has developed a new drug that requires very strict control on the amount of the drug placed in each capsule. It can be hazardous if too much drug is used and ineffective if not enough is used. The maximum error of estimate for the mean is 2 mg. The standard deviation is believed to be 7 mg. If 99 % confidence is desired, what size sample should be taken to be certain that the mean amount of the drug per capsule is correct? Show all your works. [2 marks]

Continued...

QUESTION 5 (Cont')

c. It is thought that stress may increase susceptibility to illness through suppression of the immune system. In an experiment to investigate this theory, 16 rats were randomly allocated to four treatment groups: no stress, mild stress, moderate stress and high stress. The stress conditions involved various amounts of restraint and electric shock. The concentration of lymphocytes (10^6 cells/ml) in the peripheral blood was measured for each rat with the results given in the following table. Calculations based on the raw data yielded SSA/SS(between) = 89.036 and SSE/SS(within) = 340.24. Test the null hypothesis at $\alpha = 0.05$.
Show all your works. [4 marks]

Source	*df*	SS	MS
SSA	3	89.036	29.679
SSE	12	340.24	28.353
SST	15		

Continued...

Appendix:

Formula

$$\hat{OR} = \frac{ad}{bc} \qquad \hat{RR} = \frac{a/(a+b)}{c/(c+d)}$$

$$P(A|B) = \frac{P(A \cap B)}{P(B)}$$

$$SE_{\bar{x}} = \sigma_{\bar{x}} = \frac{\sigma}{\sqrt{n}} \qquad n = \frac{z^2\sigma^2}{d^2} \qquad \bar{x} \pm t_{\alpha/2,n-1}\frac{s}{\sqrt{n}}$$

$$S_{XX} = \sum x_i^2 - \frac{(\sum x_i)^2}{n} \qquad S_{YY} = \sum y_i^2 - \frac{(\sum y_i)^2}{n}$$

$$S_{XY} = \sum x_i y_i - \frac{(\sum x_i)(\sum y_i)}{n}$$

$$b = \frac{S_{XY}}{S_{XX}} \qquad a = \bar{y} - b\,\bar{x} \qquad r = \frac{S_{XY}}{\sqrt{S_{XX}}\sqrt{S_{YY}}}$$

$$SSE = \Sigma(y - \hat{y})^2 \qquad S_{Y|X} = \sqrt{\frac{\Sigma(y_i - \hat{y}_i)^2}{n-2}}$$

$$X^2 = \sum_{all\ cells} \frac{(f_0 - f_e)^2}{f_e}$$

NEGATIVE *z* Scores



TABLE A-2 Standard Normal (z) Distribution: Cumulative Area from the LEFT

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
−3.50 and lower	.0001									
−3.4	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0003	.0002
−3.3	.0005	.0005	.0005	.0004	.0004	.0004	.0004	.0004	.0004	.0003
−3.2	.0007	.0007	.0006	.0006	.0006	.0006	.0006	.0005	.0005	.0005
−3.1	.0010	.0009	.0009	.0009	.0008	.0008	.0008	.0008	.0007	.0007
−3.0	.0013	.0013	.0013	.0012	.0012	.0011	.0011	.0011	.0010	.0010
−2.9	.0019	.0018	.0018	.0017	.0016	.0016	.0015	.0015	.0014	.0014
−2.8	.0026	.0025	.0024	.0023	.0023	.0022	.0021	.0021	.0020	.0019
−2.7	.0035	.0034	.0033	.0032	.0031	.0030	.0029	.0028	.0027	.0026
−2.6	.0047	.0045	.0044	.0043	.0041	.0040	.0039	.0038	.0037	.0036
−2.5	.0062	.0060	.0059	.0057	.0055	.0054	.0052	.0051 *	.0049	.0048
−2.4	.0082	.0080	.0078	.0075	.0073	.0071	.0069	.0068	.0066	.0064
−2.3	.0107	.0104	.0102	.0099	.0096	.0094	.0091	.0089	.0087	.0084
−2.2	.0139	.0136	.0132	.0129	.0125	.0122	.0119	.0116	.0113	.0110
−2.1	.0179	.0174	.0170	.0166	.0162	.0158	.0154	.0150	.0146	.0143
−2.0	.0228	.0222	.0217	.0212	.0207	.0202	.0197	.0192	.0188	.0183
−1.9	.0287	.0281	.0274	.0268	.0262	.0256	.0250	.0244	.0239	.0233
−1.8	.0359	.0351	.0344	.0336	.0329	.0322	.0314	.0307	.0301	.0294
−1.7	.0446	.0436	.0427	.0418	.0409	.0401	.0392	.0384	.0375	.0367
−1.6	.0548	.0537	.0526	.0516	.0505 *	.0495	.0485	.0475	.0465	.0455
−1.5	.0668	.0655	.0643	.0630	.0618 *	.0606	.0594	.0582	.0571	.0559
−1.4	.0808	.0793	.0778	.0764	.0749	.0735	.0721	.0708	.0694	.0681
−1.3	.0968	.0951	.0934	.0918	.0901	.0885	.0869	.0853	.0838	.0823
−1.2	.1151	.1131	.1112	.1093	.1075	.1056	.1038	.1020	.1003	.0985
−1.1	.1357	.1335	.1314	.1292	.1271	.1251	.1230	.1210	.1190	.1170
−1.0	.1587	.1562	.1539	.1515	.1492	.1469	.1446	.1423	.1401	.1379
−0.9	.1841	.1814	.1788	.1762	.1736	.1711	.1685	.1660	.1635	.1611
−0.8	.2119	.2090	.2061	.2033	.2005	.1977	.1949	.1922	.1894	.1867
−0.7	.2420	.2389	.2358	.2327	.2296	.2266	.2236	.2206	.2177	.2148
−0.6	.2743	.2709	.2676	.2643	.2611	.2578	.2546	.2514	.2483	.2451
−0.5	.3085	.3050	.3015	.2981	.2946	.2912	.2877	.2843	.2810	.2776
−0.4	.3446	.3409	.3372	.3336	.3300	.3264	.3228	.3192	.3156	.3121
−0.3	.3821	.3783	.3745	.3707	.3669	.3632	.3594	.3557	.3520	.3483
−0.2	.4207	.4168	.4129	.4090	.4052	.4013	.3974	.3936	.3897	.3859
−0.1	.4602	.4562	.4522	.4483	.4443	.4404	.4364	.4325	.4286	.4247
−0.0	.5000	.4960	.4920	.4880	.4840	.4801	.4761	.4721	.4681	.4641

NOTE: For values of z below −3.49, use 0.0001 for the area.

*Use these common values that result from interpolation:

z score	Area
−1.645	0.0500
−2.575	0.0050



POSITIVE z Scores

TABLE A-2 (*continued*) Cumulative Area from the LEFT

z	.00	.01	.02	.03	.04	.05	.06	.07	.08	.09
0.0	.5000	.5040	.5080	.5120	.5160	.5199	.5239	.5279	.5319	.5359
0.1	.5398	.5438	.5478	.5517	.5557	.5596	.5636	.5675	.5714	.5753
0.2	.5793	.5832	.5871	.5910	.5948	.5987	.6026	.6064	.6103	.6141
0.3	.6179	.6217	.6255	.6293	.6331	.6368	.6406	.6443	.6480	.6517
0.4	.6554	.6591	.6628	.6664	.6700	.6736	.6772	.6808	.6844	.6879
0.5	.6915	.6950	.6985	.7019	.7054	.7088	.7123	.7157	.7190	.7224
0.6	.7257	.7291	.7324	.7357	.7389	.7422	.7454	.7486	.7517	.7549
0.7	.7580	.7611	.7642	.7673	.7704	.7734	.7764	.7794	.7823	.7852
0.8	.7881	.7910	.7939	.7967	.7995	.8023	.8051	.8078	.8106	.8133
0.9	.8159	.8186	.8212	.8238	.8264	.8289	.8315	.8340	.8365	.8389
1.0	.8413	.8438	.8461	.8485	.8508	.8531	.8554	.8577	.8599	.8621
1.1	.8643	.8665	.8686	.8708	.8729	.8749	.8770	.8790	.8810	.8830
1.2	.8849	.8869	.8888	.8907	.8925	.8944	.8962	.8980	.8997	.9015
1.3	.9032	.9049	.9066	.9082	.9099	.9115	.9131	.9147	.9162	.9177
1.4	.9192	.9207	.9222	.9236	.9251	.9265	.9279	.9292	.9306	.9319
1.5	.9332	.9345	.9357	.9370	.9382	.9394	.9406	.9418	.9429	.9441
1.6	.9452	.9463	.9474	.9484	.9495 *	.9505	.9515	.9525	.9535	.9545
1.7	.9554	.9564	.9573	.9582	.9591	.9599	.9608	.9616	.9625	.9633
1.8	.9641	.9649	.9656	.9664	.9671	.9678	.9686	.9693	.9699	.9706
1.9	.9713	.9719	.9726	.9732	.9738	.9744	.9750	.9756	.9761	.9767
2.0	.9772	.9778	.9783	.9788	.9793	.9798	.9803	.9808	.9812	.9817
2.1	.9821	.9826	.9830	.9834	.9838	.9842	.9846	.9850	.9854	.9857
2.2	.9861	.9864	.9868	.9871	.9875	.9878	.9881	.9884	.9887	.9890
2.3	.9893	.9896	.9898	.9901	.9904	.9906	.9909	.9911	.9913	.9916
2.4	.9918	.9920	.9922	.9925	.9927	.9929	.9931	.9932	.9934	.9936
2.5	.9938	.9940	.9941	.9943	.9945	.9946	.9948	.9949 *	.9951	.9952
2.6	.9953	.9955	.9956	.9957	.9959	.9960	.9961	.9962	.9963	.9964
2.7	.9965	.9966	.9967	.9968	.9969	.9970	.9971	.9972	.9973	.9974
2.8	.9974	.9975	.9976	.9977	.9977	.9978	.9979	.9979	.9980	.9981
2.9	.9981	.9982	.9982	.9983	.9984	.9984	.9985	.9985	.9986	.9986
3.0	.9987	.9987	.9987	.9988	.9988	.9989	.9989	.9989	.9990	.9990
3.1	.9990	.9991	.9991	.9991	.9992	.9992	.9992	.9992	.9993	.9993
3.2	.9993	.9993	.9994	.9994	.9994	.9994	.9994	.9995	.9995	.9995
3.3	.9995	.9995	.9995	.9996	.9996	.9996	.9996	.9996	.9996	.9997
3.4	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9997	.9998
3.50 and up	.9999									

NOTE: For values of z above 3.49, use 0.9999 for the area.
*Use these common values that result from interpolation:

z score	Area
1.645	0.9500
2.575	0.9950

Common Critical Values

Confidence Level	Critical Value
0.90	1.645
0.95	1.96
0.99	2.575

TABLE A-3 t Distribution: Critical t Values

Degrees of Freedom	Area in One Tail: 0.005	0.01	0.025	0.05	0.10
	Area in Two Tails: 0.01	0.02	0.05	0.10	0.20
1	**63.657**	31.821	**12.706**	6.314	3.078
2	**9.925**	6.965	**4.303**	2.920	1.886
3	**5.841**	4.541	**3.182**	2.353	1.638
4	**4.604**	3.747	**2.776**	2.132	1.533
5	**4.032**	3.365	**2.571**	2.015	1.476
6	**3.707**	3.143	**2.447**	1.943	1.440
7	**3.499**	2.998	**2.365**	1.895	1.415
8	**3.355**	2.896	**2.306**	1.860	1.397
9	**3.250**	2.821	**2.262**	1.833	1.383
10	**3.169**	2.764	**2.228**	1.812	1.372
11	**3.106**	2.718	**2.201**	1.796	1.363
12	**3.055**	2.681	**2.179**	1.782	1.356
13	**3.012**	2.650	**2.160**	1.771	1.350
14	**2.977**	2.624	**2.145**	1.761	1.345
15	**2.947**	2.602	**2.131**	1.753	1.341
16	**2.921**	2.583	**2.120**	1.746	1.337
17	**2.898**	2.567	**2.110**	1.740	1.333
18	**2.878**	2.552	**2.101**	1.734	1.330
19	**2.861**	2.539	**2.093**	1.729	1.328
20	**2.845**	2.528	**2.086**	1.725	1.325
21	**2.831**	2.518	**2.080**	1.721	1.323
22	**2.819**	2.508	**2.074**	1.717	1.321
23	**2.807**	2.500	**2.069**	1.714	1.319
24	**2.797**	2.492	**2.064**	1.711	1.318
25	**2.787**	2.485	**2.060**	1.708	1.316
26	**2.779**	2.479	**2.056**	1.706	1.315
27	**2.771**	2.473	**2.052**	1.703	1.314
28	**2.763**	2.467	**2.048**	1.701	1.313
29	**2.756**	2.462	**2.045**	1.699	1.311
30	**2.750**	2.457	**2.042**	1.697	1.310
31	**2.744**	2.453	**2.040**	1.696	1.309
32	**2.738**	2.449	**2.037**	1.694	1.309
34	**2.728**	2.441	**2.032**	1.691	1.307
36	**2.719**	2.434	**2.028**	1.688	1.306
38	**2.712**	2.429	**2.024**	1.686	1.304
40	**2.704**	2.423	**2.021**	1.684	1.303
45	**2.690**	2.412	**2.014**	1.679	1.301
50	**2.678**	2.403	**2.009**	1.676	1.299
55	**2.668**	2.396	**2.004**	1.673	1.297
60	**2.660**	2.390	**2.000**	1.671	1.296
65	**2.654**	2.385	**1.997**	1.669	1.295
70	**2.648**	2.381	**1.994**	1.667	1.294
75	**2.643**	2.377	**1.992**	1.665	1.293
80	**2.639**	2.374	**1.990**	1.664	1.292
90	**2.632**	2.368	**1.987**	1.662	1.291
100	**2.626**	2.364	**1.984**	1.660	1.290
200	**2.601**	2.345	**1.972**	1.653	1.286
300	**2.592**	2.339	**1.968**	1.650	1.284
400	**2.588**	2.336	**1.966**	1.649	1.284
500	**2.586**	2.334	**1.965**	1.648	1.283
750	**2.582**	2.331	**1.963**	1.647	1.283
1000	**2.581**	2.330	**1.962**	1.646	1.282
2000	**2.578**	2.328	**1.961**	1.646	1.282
Large	**2.576**	2.326	**1.960**	1.645	1.282







TABLE A-4 Chi-Square (χ^2) Distribution

Degrees of Freedom	Area to the Right of the Critical Value									
	0.995	0.99	0.975	0.95	0.90	0.10	0.05	0.025	0.01	0.005
1	—	—	0.001	0.004	0.016	2.706	3.841	5.024	6.635	7.879
2	0.010	0.020	0.051	0.103	0.211	4.605	5.991	7.378	9.210	10.597
3	0.072	0.115	0.216	0.352	0.584	6.251	7.815	9.348	11.345	12.838
4	0.207	0.297	0.484	0.711	1.064	7.779	9.488	11.143	13.277	14.860
5	0.412	0.554	0.831	1.145	1.610	9.236	11.071	12.833	15.086	16.750
6	0.676	0.872	1.237	1.635	2.204	10.645	12.592	14.449	16.812	18.548
7	0.989	1.239	1.690	2.167	2.833	12.017	14.067	16.013	18.475	20.278
8	1.344	1.646	2.180	2.733	3.490	13.362	15.507	17.535	20.090	21.955
9	1.735	2.088	2.700	3.325	4.168	14.684	16.919	19.023	21.666	23.589
10	2.156	2.558	3.247	3.940	4.865	15.987	18.307	20.483	23.209	25.188
11	2.603	3.053	3.816	4.575	5.578	17.275	19.675	21.920	24.725	26.757
12	3.074	3.571	4.404	5.226	6.304	18.549	21.026	23.337	26.217	28.299
13	3.565	4.107	5.009	5.892	7.042	19.812	22.362	24.736	27.688	29.819
14	4.075	4.660	5.629	6.571	7.790	21.064	23.685	26.119	29.141	31.319
15	4.601	5.229	6.262	7.261	8.547	22.307	24.996	27.488	30.578	32.801
16	5.142	5.812	6.908	7.962	9.312	23.542	26.296	28.845	32.000	34.267
17	5.697	6.408	7.564	8.672	10.085	24.769	27.587	30.191	33.409	35.718
18	6.265	7.015	8.231	9.390	10.865	25.989	28.869	31.526	34.805	37.156
19	6.844	7.633	8.907	10.117	11.651	27.204	30.144	32.852	36.191	38.582
20	7.434	8.260	9.591	10.851	12.443	28.412	31.410	34.170	37.566	39.997
21	8.034	8.897	10.283	11.591	13.240	29.615	32.671	35.479	38.932	41.401
22	8.643	9.542	10.982	12.338	14.042	30.813	33.924	36.781	40.289	42.796
23	9.260	10.196	11.689	13.091	14.848	32.007	35.172	38.076	41.638	44.181
24	9.886	10.856	12.401	13.848	15.659	33.196	36.415	39.364	42.980	45.559
25	10.520	11.524	13.120	14.611	16.473	34.382	37.652	40.646	44.314	46.928
26	11.160	12.198	13.844	15.379	17.292	35.563	38.885	41.923	45.642	48.290
27	11.808	12.879	14.573	16.151	18.114	36.741	40.113	43.194	46.963	49.645
28	12.461	13.565	15.308	16.928	18.939	37.916	41.337	44.461	48.278	50.993
29	13.121	14.257	16.047	17.708	19.768	39.087	42.557	45.722	49.588	52.336
30	13.787	14.954	16.791	18.493	20.599	40.256	43.773	46.979	50.892	53.672
40	20.707	22.164	24.433	26.509	29.051	51.805	55.758	59.342	63.691	66.766
50	27.991	29.707	32.357	34.764	37.689	63.167	67.505	71.420	76.154	79.490
60	35.534	37.485	40.482	43.188	46.459	74.397	79.082	83.298	88.379	91.952
70	43.275	45.442	48.758	51.739	55.329	85.527	90.531	95.023	100.425	104.215
80	51.172	53.540	57.153	60.391	64.278	96.578	101.879	106.629	112.329	116.321
90	59.196	61.754	65.647	69.126	73.291	107.565	113.145	118.136	124.116	128.299
100	67.328	70.065	74.222	77.929	82.358	118.498	124.342	129.561	135.807	140.169

From Donald B. Owen, *Handbook of Statistical Tables*, © 1962 Addison-Wesley Publishing Co., Reading, MA. Reprinted with permission of the publisher.

Degrees of Freedom

$n - 1$	for confidence intervals or hypothesis tests with a standard deviation or variance
$k - 1$	for multinomial experiments or goodness-of-fit with k categories
$(r - 1)(c - 1)$	for contingency tables with r rows and c columns
$k - 1$	for Kruskal-Wallis test with k samples

TABLE A-5 *F* Distribution ($\alpha = 0.025$ in the right tail)

Denominator degrees of freedom (df_2)	Numerator degrees of freedom (df_1)									
	1	2	3	4	5	6	7	8	9
1	647.79	799.50	864.16	899.58	921.85	937.11	948.22	956.66	963.28
2	38.506	39.000	39.165	39.248	39.298	39.331	39.355	39.373	39.387
3	17.443	16.044	15.439	15.101	14.885	14.735	14.624	14.540	14.473
4	12.218	10.649	9.9792	9.6045	9.3645	9.1973	9.0741	8.9796	8.9047
5	10.007	8.4336	7.7636	7.3879	7.1464	6.9777	6.8531	6.7572	6.6811
6	8.8131	7.2599	6.5988	6.2272	5.9876	5.8198	5.6955	5.5996	5.5234
7	8.0727	6.5415	5.8898	5.5226	5.2852	5.1186	4.9949	4.8993	4.8232
8	7.5709	6.0595	5.4160	5.0526	4.8173	4.6517	4.5286	4.4333	4.3572
9	7.2093	5.7147	5.0781	4.7181	4.4844	4.3197	4.1970	4.1020	4.0260
10	6.9367	5.4564	4.8256	4.4683	4.2361	4.0721	3.9498	3.8549	3.7790
11	6.7241	5.2559	4.6300	4.2751	4.0440	3.8807	3.7586	3.6638	3.5879
12	6.5538	5.0959	4.4742	4.1212	3.8911	3.7283	3.6065	3.5118	3.4358
13	6.4143	4.9653	4.3472	3.9959	3.7667	3.6043	3.4827	3.3880	3.3120
14	6.2979	4.8567	4.2417	3.8919	3.6634	3.5014	3.3799	3.2853	3.2093
15	6.1995	4.7650	4.1528	3.8043	3.5764	3.4147	3.2934	3.1987	3.1227
16	6.1151	4.6867	4.0768	3.7294	3.5021	3.3406	3.2194	3.1248	3.0488
17	6.0420	4.6189	4.0112	3.6648	3.4379	3.2767	3.1556	3.0610	2.9849
18	5.9781	4.5597	3.9539	3.6083	3.3820	3.2209	3.0999	3.0053	2.9291
19	5.9216	4.5075	3.9034	3.5587	3.3327	3.1718	3.0509	2.9563	2.8801
20	5.8715	4.4613	3.8587	3.5147	3.2891	3.1283	3.0074	2.9128	2.8365
21	5.8266	4.4199	3.8188	3.4754	3.2501	3.0895	2.9686	2.8740	2.7977
22	5.7863	4.3828	3.7829	3.4401	3.2151	3.0546	2.9338	2.8392	2.7628
23	5.7498	4.3492	3.7505	3.4083	3.1835	3.0232	2.9023	2.8077	2.7313
24	5.7166	4.3187	3.7211	3.3794	3.1548	2.9946	2.8738	2.7791	2.7027
25	5.6864	4.2909	3.6943	3.3530	3.1287	2.9685	2.8478	2.7531	2.6766
26	5.6586	4.2655	3.6697	3.3289	3.1048	2.9447	2.8240	2.7293	2.6528
27	5.6331	4.2421	3.6472	3.3067	3.0828	2.9228	2.8021	2.7074	2.6309
28	5.6096	4.2205	3.6264	3.2863	3.0626	3.9027	2.7820	2.6872	2.6106
29	5.5878	4.2006	3.6072	3.2674	3.0438	2.8840	2.7633	2.6686	2.5919
30	5.5675	4.1821	3.5894	3.2499	3.0265	2.8667	2.7460	2.6513	2.5746
40	5.4239	4.0510	3.4633	3.1261	2.9037	2.7444	2.6238	2.5289	2.4519
60	5.2856	3.9253	3.3425	3.0077	2.7863	2.6274	2.5068	2.4117	2.3344
120	5.1523	3.8046	3.2269	2.8943	2.6740	2.5154	2.3948	2.2994	2.2217
∞	5.0239	3.6889	3.1161	2.7858	2.5665	2.4082	2.2875	2.1918	2.1136

TABLE A-5 *F* Distribution ($\alpha = 0.025$ in the right tail) *(continued)*

Denominator degrees of freedom (df_2) \ Numerator degrees of freedom (df_1)	10	12	15	20	24	30	40	60	120	∞
1	968.63	976.71	984.87	993.10	997.25	1001.4	1005.6	1009.8	1014.0	1018.3
2	39.398	39.415	39.431	39.448	39.456	39.465	39.473	39.481	39.490	39.498
3	14.419	14.337	14.253	14.167	14.124	14.081	14.037	13.992	13.947	13.902
4	8.8439	8.7512	8.6565	8.5599	8.5109	8.4613	8.4111	8.3604	8.3092	8.2573
5	6.6192	6.5245	6.4277	6.3286	6.2780	6.2269	6.1750	6.1225	6.0693	6.0153
6	5.4613	5.3662	5.2687	5.1684	5.1172	5.0652	5.0125	4.9589	4.9044	4.8491
7	4.7611	4.6658	4.5678	4.4667	4.4150	4.3624	4.3089	4.2544	4.1989	4.1423
8	4.2951	4.1997	4.1012	3.9995	3.9472	3.8940	3.8398	3.7844	3.7279	3.6702
9	3.9639	3.8682	3.7694	3.6669	3.6142	3.5604	3.5055	3.4493	3.3918	3.3329
10	3.7168	3.6209	3.5217	3.4185	3.3654	3.3110	3.2554	3.1984	3.1399	3.0798
11	3.5257	3.4296	3.3299	3.2261	3.1725	3.1176	3.0613	3.0035	2.9441	2.8828
12	3.3736	3.2773	3.1772	3.0728	3.0187	2.9633	2.9063	2.8478	2.7874	2.7249
13	3.2497	3.1532	3.0527	2.9477	2.8932	2.8372	2.7797	2.7204	2.6590	2.5955
14	3.1469	3.0502	2.9493	2.8437	2.7888	2.7324	2.6742	2.6142	2.5519	2.4872
15	3.0602	2.9633	2.8621	2.7559	2.7006	2.6437	2.5850	2.5242	2.4611	2.3953
16	2.9862	2.8890	2.7875	2.6808	2.6252	2.5678	2.5085	2.4471	2.3831	2.3163
17	2.9222	2.8249	2.7230	2.6158	2.5598	2.5020	2.4422	2.3801	2.3153	2.2474
18	2.8664	2.7689	2.6667	2.5590	2.5027	2.4445	2.3842	2.3214	2.2558	2.1869
19	2.8172	2.7196	2.6171	2.5089	2.4523	2.3937	2.3329	2.2696	2.2032	2.1333
20	2.7737	2.6758	2.5731	2.4645	2.4076	2.3486	2.2873	2.2234	2.1562	2.0853
21	2.7348	2.6368	2.5338	2.4247	2.3675	2.3082	2.2465	2.1819	2.1141	2.0422
22	2.6998	2.6017	2.4984	2.3890	2.3315	2.2718	2.2097	2.1446	2.0760	2.0032
23	2.6682	2.5699	2.4665	2.3567	2.2989	2.2389	2.1763	2.1107	2.0415	1.9677
24	2.6396	2.5411	2.4374	2.3273	2.2693	2.2090	2.1460	2.0799	2.0099	1.9353
25	2.6135	2.5149	2.4110	2.3005	2.2422	2.1816	2.1183	2.0516	1.9811	1.9055
26	2.5896	2.4908	2.3867	2.2759	2.2174	2.1565	2.0928	2.0257	1.9545	1.8781
27	2.5676	2.4688	2.3644	2.2533	2.1946	2.1334	2.0693	2.0018	1.9299	1.8527
28	2.5473	2.4484	2.3438	2.2324	2.1735	2.1121	2.0477	1.9797	1.9072	1.8291
29	2.5286	2.4295	2.3248	2.2131	2.1540	2.0923	2.0276	1.9591	1.8861	1.8072
30	2.5112	2.4120	2.3072	2.1952	2.1359	2.0739	2.0089	1.9400	1.8664	1.7867
40	2.3882	2.2882	2.1819	2.0677	2.0069	1.9429	1.8752	1.8028	1.7242	1.6371
60	2.2702	2.1692	2.0613	1.9445	1.8817	1.8152	1.7440	1.6668	1.5810	1.4821
120	2.1570	2.0548	1.9450	1.8249	1.7597	1.6899	1.6141	1.5299	1.4327	1.3104
∞	2.0483	1.9447	1.8326	1.7085	1.6402	1.5660	1.4835	1.3883	1.2684	1.0000

From Maxine Merrington and Catherine M. Thompson, "Tables of Percentage Points of the Inverted Beta (*F*) Distribution," *Biometrika 33* (1943): 80–84. Reproduced with permission of the Biometrika Trustees.

(continued)

TABLE A-5 *F* Distribution ($\alpha = 0.05$ in the right tail)

Denominator degrees of freedom (df_2)	Numerator degrees of freedom (df_1)								
	1	2	3	4	5	6	7	8	9
1	161.45	199.50	215.71	224.58	230.16	233.99	236.77	238.88	240.54
2	18.513	19.000	19.164	19.247	19.296	19.330	19.353	19.371	19.385
3	10.128	9.5521	9.2766	9.1172	9.0135	8.9406	8.8867	8.8452	8.8123
4	7.7086	6.9443	6.5914	6.3882	6.2561	6.1631	6.0942	6.0410	6.9988
5	6.6079	5.7861	5.4095	5.1922	5.0503	4.9503	4.8759	4.8183	4.7725
6	5.9874	5.1433	4.7571	4.5337	4.3874	4.2839	4.2067	4.1468	4.0990
7	5.5914	4.7374	4.3468	4.1203	3.9715	3.8660	3.7870	3.7257	3.6767
8	5.3177	4.4590	4.0662	3.8379	3.6875	3.5806	3.5005	3.4381	3.3881
9	5.1174	4.2565	3.8625	3.6331	3.4817	3.3738	3.2927	3.2296	3.1789
10	4.9646	4.1028	3.7083	3.4780	3.3258	3.2172	3.1355	3.0717	3.0204
11	4.8443	3.9823	3.5874	3.3567	3.2039	3.0946	3.0123	2.9480	2.8962
12	4.7472	3.8853	3.4903	3.2592	3.1059	2.9961	2.9134	2.8486	2.7964
13	4.6672	3.8056	3.4105	3.1791	3.0254	2.9153	2.8321	2.7669	2.7144
14	4.6001	3.7389	3.3439	3.1122	2.9582	2.8477	2.7642	2.6987	2.6458
15	4.5431	3.6823	3.2874	3.0556	2.9013	2.7905	2.7066	2.6408	2.5876
16	4.4940	3.6337	3.2389	3.0069	2.8524	2.7413	2.6572	2.5911	2.5377
17	4.4513	3.5915	3.1968	2.9647	2.8100	2.6987	2.6143	2.5480	2.4943
18	4.4139	3.5546	3.1599	2.9277	2.7729	2.6613	2.5767	2.5102	2.4563
19	4.3807	3.5219	3.1274	2.8951	2.7401	2.6283	2.5435	2.4768	2.4227
20	4.3512	3.4928	3.0984	2.8661	2.7109	2.5990	2.5140	2.4471	2.3928
21	4.3248	3.4668	3.0725	2.8401	2.6848	2.5727	2.4876	2.4205	2.3660
22	4.3009	3.4434	3.0491	2.8167	2.6613	2.5491	2.4638	2.3965	2.3419
23	4.2793	3.4221	3.0280	2.7955	2.6400	2.5277	2.4422	2.3748	2.3201
24	4.2597	3.4028	3.0088	2.7763	2.6207	2.5082	2.4226	2.3551	2.3002
25	4.2417	3.3852	2.9912	2.7587	2.6030	2.4904	2.4047	2.3371	2.2821
26	4.2252	3.3690	2.9752	2.7426	2.5868	2.4741	2.3883	2.3205	2.2655
27	4.2100	3.3541	2.9604	2.7278	2.5719	2.4591	2.3732	2.3053	2.2501
28	4.1960	3.3404	2.9467	2.7141	2.5581	2.4453	2.3593	2.2913	2.2360
29	4.1830	3.3277	2.9340	2.7014	2.5454	2.4324	2.3463	2.2783	2.2229
30	4.1709	3.3158	2.9223	2.6896	2.5336	2.4205	2.3343	2.2662	2.2107
40	4.0847	3.2317	2.8387	2.6060	2.4495	2.3359	2.2490	2.1802	2.1240
60	4.0012	3.1504	2.7581	2.5252	2.3683	2.2541	2.1665	2.0970	2.0401
120	3.9201	3.0718	2.6802	2.4472	2.2899	2.1750	2.0868	2.0164	1.9588
∞	3.8415	2.9957	2.6049	2.3719	2.2141	2.0986	2.0096	1.9384	1.8799

(continued)

TABLE A-5 *F* Distribution ($\alpha = 0.05$ in the right tail) *(continued)*

Denominator degrees of freedom (df_2)	Numerator degrees of freedom (df_1)											
	10	12	15	20	24	30	40	60	120	∞
1	241.88	243.91	245.95	248.01	249.05	250.10	251.14	252.20	253.25	254.31
2	19.396	19.413	19.429	19.446	19.454	19.462	19.471	19.479	19.487	19.496
3	8.7855	8.7446	8.7029	8.6602	8.6385	8.6166	8.5944	8.5720	8.5494	8.5264
4	5.9644	5.9117	5.8578	5.8025	5.7744	5.7459	5.7170	5.6877	5.6581	5.6281
5	4.7351	4.6777	4.6188	4.5581	4.5272	4.4957	4.4638	4.4314	4.3985	4.3650
6	4.0600	3.9999	3.9381	3.8742	3.8415	3.8082	3.7743	3.7398	3.7047	3.6689
7	3.6365	3.5747	3.5107	3.4445	3.4105	3.3758	3.3404	3.3043	3.2674	3.2298
8	3.3472	3.2839	3.2184	3.1503	3.1152	3.0794	3.0428	3.0053	2.9669	2.9276
9	3.1373	3.0729	3.0061	2.9365	2.9005	2.8637	2.8259	2.7872	2.7475	2.7067
10	2.9782	2.9130	2.8450	2.7740	2.7372	2.6996	2.6609	2.6211	2.5801	2.5379
11	2.8536	2.7876	2.7186	2.6464	2.6090	2.5705	2.5309	2.4901	2.4480	2.4045
12	2.7534	2.6866	2.6169	2.5436	2.5055	2.4663	2.4259	2.3842	2.3410	2.2962
13	2.6710	2.6037	2.5331	2.4589	2.4202	2.3803	2.3392	2.2966	2.2524	2.2064
14	2.6022	2.5342	2.4630	2.3879	2.3487	2.3082	2.2664	2.2229	2.1778	2.1307
15	2.5437	2.4753	2.4034	2.3275	2.2878	2.2468	2.2043	2.1601	2.1141	2.0658
16	2.4935	2.4247	2.3522	2.2756	2.2354	2.1938	2.1507	2.1058	2.0589	2.0096
17	2.4499	2.3807	2.3077	2.2304	2.1898	2.1477	2.1040	2.0584	2.0107	1.9604
18	2.4117	2.3421	2.2686	2.1906	2.1497	2.1071	2.0629	2.0166	1.9681	1.9168
19	2.3779	2.3080	2.2341	2.1555	2.1141	2.0712	2.0264	1.9795	1.9302	1.8780
20	2.3479	2.2776	2.2033	2.1242	2.0825	2.0391	1.9938	1.9464	1.8963	1.8432
21	2.3210	2.2504	2.1757	2.0960	2.0540	2.0102	1.9645	1.9165	1.8657	1.8117
22	2.2967	2.2258	2.1508	2.0707	2.0283	1.9842	1.9380	1.8894	1.8380	1.7831
23	2.2747	2.2036	2.1282	2.0476	2.0050	1.9605	1.9139	1.8648	1.8128	1.7570
24	2.2547	2.1834	2.1077	2.0267	1.9838	1.9390	1.8920	1.8424	1.7896	1.7330
25	2.2365	2.1649	2.0889	2.0075	1.9643	1.9192	1.8718	1.8217	1.7684	1.7110
26	2.2197	2.1479	2.0716	1.9898	1.9464	1.9010	1.8533	1.8027	1.7488	1.6906
27	2.2043	2.1323	2.0558	1.9736	1.9299	1.8842	1.8361	1.7851	1.7306	1.6717
28	2.1900	2.1179	2.0411	1.9586	1.9147	1.8687	1.8203	1.7689	1.7138	1.6541
29	2.1768	2.1045	2.0275	1.9446	1.9005	1.8543	1.8055	1.7537	1.6981	1.6376
30	2.1646	2.0921	2.0148	1.9317	1.8874	1.8409	1.7918	1.7396	1.6835	1.6223
40	2.0772	2.0035	1.9245	1.8389	1.7929	1.7444	1.6928	1.6373	1.5766	1.5089
60	1.9926	1.9174	1.8364	1.7480	1.7001	1.6491	1.5943	1.5343	1.4673	1.3893
120	1.9105	1.8337	1.7505	1.6587	1.6084	1.5543	1.4952	1.4290	1.3519	1.2539
∞	1.8307	1.7522	1.6664	1.5705	1.5173	1.4591	1.3940	1.3180	1.2214	1.0000

From Maxine Merrington and Catherine M. Thompson. "Tables of Percentage Points of the Inverted Beta (*F*) Distribution," *Biometrika 33* (1943): 80–84. Reproduced with permission of the Biometrika Trustees.

End of Paper